# DEPARTMENT of ENVIRONMENTAL SERVICES Water Division - Watershed Management Bureau

#### LAKE TROPHIC DATA

## MORPHOMETRIC:

Lake: TRIO PONDS, ONE AND TWO 27.44 Lake Area (ha): Maximum depth (m): Town: ODELL 10.6 Mean depth (m): County: Coos 3.7 River Basin: Connecticut Volume (m<sup>3</sup>): 1002500 Latitude: 44°42'38" N Longitude: 71°22'00" W Relative depth: 1.8 Shore configuration: 1.83 Elevation (ft): 2315
Shore length (m): 3400
Watershed area (ha): 338.4 Areal water load (m/yr): 10.84 Flushing rate (yr<sup>-1</sup>): 3.00 P retention coeff.: 0.54 % watershed ponded: 2.3 Lake type: natural w/dam

BIOLOGICAL:	8 March 1999	3 August 1998
DOM. PHYTOPLANKTON (% TOTAL) #1	NO WINTER PLANKTON	DINOBRYON 40%
#2	ANALYZED	ASTERIONELLA 25%
#3		MERISMOPEDIA 6%
PHYTOPLANKTON ABUNDANCE (units/mL)		
CHLOROPHYLL-A (µg/L)		2.37
DOM. ZOOPLANKTON (% TOTAL) #1		ROTIFER SPP 41%
#2		NAUPLIUS LARVA 12%
#3		
ROTIFERS/LITER		36
MICROCRUSTACEA/LITER		27
ZOOPLANKTON ABUNDANCE (#/L)		73
VASCULAR PLANT ABUNDANCE		Scattered
SECCHI DISK TRANSPARENCY (m)		3.6
BOTTOM DISSOLVED OXYGEN (mg/L)	5.5	1.5
BACTERIA (E. coli, #/100 ml) #1		
#2		
#3		

# SUMMER THERMAL STRATIFICATION:

#### stratified

Depth of thermocline (m): 4.4
Hypolimnion volume (m³): 159500
Anoxic volume (m³): None

CHEMICAL:			TRIO PONI	OS, ONE AN	OWT O
	8 March 1999		3 August 1998		
DEPTH (m)	2.5	5.0	1.5	4.0	7.5
pH (units)	5.5	5.4	6.1	5.0	5.2
A.N.C. (Alkalinity)	1.6	1.9	2.4	1.2	1.4
NITRATE NITROGEN	0.10	0.08	< 0.05	< 0.05	< 0.05
TOTAL KJELDAHL NITROGEN	0.10	0.20	0.30	0.30	0.20
TOTAL PHOSPHORUS	0.006	0.007	0.010	0.007	0.011
CONDUCTIVITY (µmhos/cm)	20.1	20.5	15.1	15.0	17.1
APPARENT COLOR (cpu)	42	42	52	55	58
MAGNESIUM			0.21		
CALCIUM			1.2		
SODIUM			< 1.0		
POTASSIUM			0.26	-	
CHLORIDE	< 2	< 2	< 2	< 2	< 2
SULFATE	4	4	3	3	3
TN : TP	33	40	30	43	18
CALCITE SATURATION INDEX			4.7		


All results in mg/L unless indicated otherwise

TROPHIC CLASSIFICATION: 1998

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
2	2	1	0	5	Oligo.

# **COMMENTS:**

- 1. aka Lower Trio Pond or Big Trio Pond.
- 2. A chain of three partially manmade ponds were built by Groveton Paper Company in 1914 for log driving purposes, and were known as Trio Ponds #1, #2, and #3. The dam on pond #1 was later raised so that #1 and #2 ponds were joined into one pond (and often called Lower Trio Pond).
- 3. This is a tea-colored, acidic to moderately acidic, relatively unproductive remote pond accessible by 4-wheel drive only.
- 4. Approximately 10 camps were located along the southwestern end of the pond near the dam.



## FIELD DATA SHEET

LAKE: TRIO PONDS, ONE AND TWO TOWN: ODELL
DATE: 08/03/1998 WEATHER: PRTLY CLOUDY & MILD; LT BREEZE


DATE: 00/03/1330	WENTIL	EK: IKIHI CHOODI W	MIED, EI BREEZE
DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION
0.1	20.0	8.6	93 %
1.0	20.0	8.6	93 %
2.0	20.0	8.5	93 %
3.0	19.0	8.2	88 %
4.0	17.0	6.5	67 %
5.0	12.0	4.2	39 %
6.0	10.8	3.8	33 %
7.0	10.5	3.5	31 %
8.0	10.2	2.8	24 %
9.0	10.0	1.5	13 %
		And the Marketon Control of the Cont	
	1		
·			

SECCHI DISK (m): 3.6 COMMENTS:

BOTTOM DEPTH (m): 9.2

TIME: 1340

\*Dissolved oxygen values are in mg/L



## AQUATIC PLANT SURVEY

TOWN: ODELL

PLANT NAME Key **ABUNDANCE GENERIC** COMMON Ε Eriocaulon septangulare Pipewort Scattered L Lobelia dortmanna Water lobelia Sparse W Potamogeton Pondweed Sparse

Sweet gale

Filamentous algae

Freshwater sponge

S Sparganium Bur reed Sparse
G Gramineae Grass family Sparse
h Hypericum St. John's-wort Sparse

 d
 Dulichium arundinaceum
 Three-way sedge
 Sparse

 Y
 Nuphar
 Yellow water lily
 Sparse

Brasenia schreberi Water shield Sparse

Vaccinium corymbosum High-bush blueberry Scattered

 N
 Nymphaea
 White water lily
 Sparse

 X
 Sterile thread-like leaf
 Sparse

bScirpusBulrushSparseUUtriculariaBladderwortSparse

D Drosera Sundew Sparse

I Iris Iris Sparse

OVERALL ABUNDANCE: Scattered

DATE: 08/03/1998

Scattered

Sparse

Sparse

# GENERAL OBSERVATIONS:

LAKE: TRIO PONDS, ONE AND TWO

Myrica gale

Spongilla

g f

a

В

V

1. A family of three loons and a hawk were observed.